

Curculigo latifolia Dryand ex W. T. Aiton

Identifiants : 121858/curclati

Association du Potager de mes (nos) Rêves
(<https://lepotager-demesreves.fr> - <https://terralisme.fr>)

Fiche réalisée par Patrick Le Ménahèze

Dernière modification le 08/08/2026

- **Classification/taxinomie traditionnelle :**

- Règne : Fungi ;
- Division : Basidiomycota ;
- Classe : Agaricomycetes ;
- Ordre : Agaricales ;
- Famille : Hypoxidaceae ;
- Genre : Curculigo ;



- **Synonymes :** Curculigo borneensis Merr.; Curculigo latifolia Dryand. ex W. T. Aiton ; Curculigo longiflora Kurz; Molineria longiflora Kurz; Molineria latifolia (Dryand ex W. T. Aiton) ; and several others ;

- **Nom(s) anglais, local(aux) et/ou international(aux) :** , Daun matahari, Dong sbat, Lamba biasa, Lemba, Lembah, Lumbah, Lunbah rimba, Lunbah padi, Kelapa puyoh, Labpak, Rumpit palem, Tambaka, ;



- **Rapport de consommation et comestibilité/consommabilité inférée (partie(s) utilisable(s) et usage(s) alimentaire(s) correspondant(s)) :**

Détails : Les fruits sont mangés. Les fruits ont un goût légèrement sucré et, une fois consommés, les autres aliments ont un goût sucré pendant environ une heure.{{{0(+x)}} (traduction automatique)

Original : The fruit are eaten. The fruit have a slightly sweet taste and after they are eaten other foods taste sweet for an hour or so{{{0(+x)}}



Précautions : néant, inconnus ou indéterminés.

- **Liens, sources et/ou références :**

dont classification :

dont biographie/références de "FOOD PLANTS INTERNATIONAL" :

Ambasta, S.P. (Ed.), 2000, The Useful Plants of India. CSIR India. p 151 (As Curculigo latifolia)Burkill, I.H., 1966, A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia. Vol 1 (A-H) p 713 (As Curculigo latifolia)Fern, K., 2012, Tropical Species Database <http://theferns.info/tropical/Hariyadi>, B., 2008, The Entwined Tree: Traditional Natural Resource Management of Serampas, Jambi, Indonesia. Ph. D thesis. Univ. of Hawaii. p 444 (As Curculigo latifolia)Hoare, A., 2003, Food use of the Lundayeh SW Sabah. Borneo Research Council. (As Curculigo latifolia)Hortus kew. ed. 2, 2:253. 1811 (As Curculigo latifolia)Milow, P., et al, 2013, Malaysian species of plants with edible fruits or seeds and their evaluation.

International Journal of Fruit Science. 14:1, 1-27 (As *Curculigo latifolia*)Natuurk. Tijdschr. Ned.-Indies 27:232. 1864Ooi, D. J., et al, 2018, Polyphenol-rich ethyl acetate fraction of *Molineria latifolia* rhizome restores oxidant-antioxidant balance by possible engagement of KEAP1-NRF2 and PKC/NF- κ B signalling pathways. Journal of Functional Foods 42 (2018) 111-121Phon, P., 2000, Plants used in Cambodia. © Pauline Dy Phon, Phnom Penh, Cambodia. p 444Rollins, C. B., 1987, The Role of the Fruit & Spice Park in Popularization and Dissemination of Tropical Fruit in South Florida. Proc. Fla. State Hort. Soc. 100:323-327 (As *Curculigo latifolia*)Singh, H.B., Arora R.K.,1978, Wild edible Plants of India. Indian Council of Agricultural Research, New Delhi. p 54 (As *Curculigo latifolia*)Sukarya, D. G., (Ed.) 2013, 3,500 Plant Species of the Botanic Gardens of Indonesia. LIPI p 1124Tropical Plants Database, Ken Fern. tropical.theferns.info.